

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/GB2006/003196

International filing date: 25 August 2006 (25.08.2006)

Document type: Certified copy of priority document

Document details: Country/Office: GB
Number: 0517386.9
Filing date: 26 August 2005 (26.08.2005)

Date of receipt at the International Bureau: 05 October 2006 (05.10.2006)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

I also certify that the attached copy of the request for grant of a Patent (Form 1/77) bears an amendment, effected by this office, following a request by the applicant and agreed to by the Comptroller-General.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.


Signed

Andrew Gersey

Dated

27 September 2006

Patents Form

Patents Act 1977
(Rule 16)J128497/ 001 D02246 P01777NDFEE....
26AUG05 0.00 ACCOUNT 0517386.9**Request for grant of a patent** 6 AUG 2005

(An explanatory leaflet on how to fill in this form is available from the Patent Office)

Application number GB**0517386.9**

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ1. Your reference:
(optional)

P023742GB JPT

2. Full name, address and postcode of the applicant or of each applicant (*underline all surnames*):Antisoma Research Limited
West Africa House
Hanger Lane
Ealing, London,
W5 3QR, United Kingdom

07869100001

Patents ADP number (*if you know it*):

If the applicant is a corporate body, give the country/state of its incorporation:

United Kingdom

3. Title of the invention:

Combinations for the treatment of cancer.

4. Name of your agent (*if you have one*):

D Young & Co

"Address for service" in the United Kingdom to which all correspondence should be sent (*including the postcode*):120 Holborn
London
EC1N 2DYPatents ADP number (*if you know it*):

59006

ERIC POTTER CLARKSON
Park View House LLP
58 The Row
Nottingham
NG1 5DD

5. Priority declaration: Are you claiming priority from one or more earlier-filed patent applications? If so, please give details of the application(s):

Country

Application number
(*if you know it*)Date of filing
(day / month / year)

6. Divisionals etc: Is this application a divisional application, or being made following resolution of an entitlement dispute about an earlier application? If so, please give the application number and filing date of the earlier application:

Number of earlier UK application

Date of filing
(day / month / year)

7. Inventorship: (Inventors must be individuals not companies)

(Please tick the appropriate boxes)

Are all the applicants named above also inventors?

YES ☐NO ☒

If yes, are there any other inventors?

YES ☐NO ☐

8. Are you paying the application fee with this form?

YES ☐NO ☒

Patents Form 1/77

Patents Form 1/77

9. Accompanying documents: not counting duplicates, please enter the number of pages of each item accompanying this form:

Continuation sheets of this form:

Description: 13

Claim (s): 7

Abstract: 1

Drawing(s): -

If you are not filing a description, please give details of the previous application you are going to rely upon:

Country

Application number

Date of filing
(day / month / year)

10. If you are also filing any of the following, state how many against each item.

Priority documents:

Statement of inventorship and right to grant of a patent (Patents Form 7/77):

Request for search (Patents Form 9A/77):

Request for substantive examination (Patents Form 10/77):

Any other documents:
(please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature(s): D. Young & Co.
D Young & Co (Agents for the Applicants)

Date: 26 August 2005

12. Name, e-mail address, telephone, Fax and/or mobile number, if any, of a contact point for the applicant:

James P Tanner

023 8071 9500

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you are resident in the United Kingdom and your application contains information which relates to military technology, or would be prejudicial to national security or the safety of the public, section 23 of the Patents Act 1977 prohibits you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

A leaflet on how to fill in this form is available from the Patent Office. If you would like a copy of the leaflet it is available on our website at <http://www.patent.gov.uk/patent/info/fact05.pdf> or alternatively you could telephone 08459 500505 or send an email to enquiries@patent.gov.uk to request a copy.


Patents Form 1/77

DUPLICATE

Combinations for the treatment of cancer

The present invention relates to combinations of compounds of the class having the formula (I) as defined below, for example compounds of the xanthenone acetic acid class having the formula (II) as defined below, such as 5,6-dimethylxanthenone-4-acetic acid (DMXAA), or a pharmaceutically acceptable salt, ester or prodrug thereof and vascular endothelial growth factor (VEGF) binders, in particular the monoclonal antibody Avastin™ (bevacizumab). For example, the present invention relates to synergistic combinations of compounds of the class having the formula (I) as defined below, for example compounds of the xanthenone acetic acid class having the formula (II) as defined below, such as 5,6-dimethylxanthenone-4-acetic acid (DMXAA), or pharmaceutically acceptable salt, ester or prodrug thereof and anti-angiogenic growth factor inhibitors, in particular the monoclonal antibody Avastin™ (bevacizumab), a VEGF binder. More particularly, the invention is concerned with the use of such combinations in the treatment of cancer. The present invention also relates to pharmaceutical compositions containing such combinations.

5,6-dimethylxanthenone-4-acetic acid (DMXAA) is represented by the following formula:



Three phase I clinical trials of DMXAA as a monotherapy have recently been completed, with dynamic MRI showing that it induces a significant reduction in tumour blood flow at well-tolerated doses. DMXAA is thus one of the first vascular disrupting agents (VDAs) for which activity (irreversible inhibition of tumour blood flow) has been documented in human tumours. These findings are in agreement with preclinical studies using syngeneic murine tumours or human tumour xenografts

which showed that its antivascular activity produced prolonged inhibition of tumour blood flow leading to extensive regions of haemorrhagic necrosis.

However, in these phase I clinical trials of DMXAA there were very few tumour responses, demonstrating that DMXAA alone does not have significant potential in cancer treatment as a single agent. Therefore, there is a need to identify compounds that could have a synergistic effect with DMXAA.

There is a new class of cancer drugs available that are not cytotoxics, but block the growth factor signalling pathways. Examples include Avastin™ (bevacizumab), a humanised monoclonal antibody that binds to vascular endothelial growth factor (VEGF). By doing so, it inhibits angiogenesis (growth of new blood vessels), starving growing tumour of nutrients. We have surprisingly found that DMXAA may act synergistically with these new agents, enhancing their anti-cancer activity.

Vascular Endothelial Growth Factor

Tumours have been found to overexpress certain growth factors that enable them to proliferate rapidly. Chief among these is VEGF. Tumours secrete VEGF, which stimulates endothelial proliferation and migration through two high-affinity receptor-associated tyrosine kinases found primarily on the vascular endothelium, VEGF-R1 (Flt-1) and VEGF-R2 (Flk-1/KDR). Expression levels of VEGF are negatively correlated with prognosis and survival in cancer, and inhibiting its binding to its receptor has been shown to improve survival.

VEGF is targeted by Avastin™ (bevacizumab, a humanised monoclonal antibody marketed by Genentech in the US and Roche elsewhere). The antibody binds directly to VEGF, preventing it from binding to VEGF receptors on the vascular endothelium. This means that the new blood vessels required by the tumour do not develop, and it cannot grow. Avastin™ combined with standard chemotherapy has been shown to offer a survival advantage over standard chemotherapy alone in colorectal, lung and breast cancers in phase III trials.

Previous DMXAA combination studies

DMXAA has previously been demonstrated to have synergy with a number of agents in xenograft studies. These agents include widely used cytotoxic chemotherapies such as taxanes (paclitaxel and docetaxel), platins (cisplatin and carboplatin), vinca alkaloids (vincristine), antimetabolites (gemcitabine), topoisomerase II inhibitors (etoposide) and anthracyclines (doxorubicin). It is believed that the synergy arises because DMXAA causes necrosis in the centre of tumours by disrupting the blood vessels that supply the core, but it leaves a viable rim of rapidly proliferating cancer cells that are supplied by normal blood vessels. These remaining malignant cells are targeted by the cytotoxic agents, which primarily act by disrupting cell division in various ways.

DMXAA is currently in two phase II trials examining its anti-tumour efficacy in combination with paclitaxel and carboplatin, and one trial combining it with docetaxel. Although the taxanes are believed to have anti-angiogenic properties, this is via a very different mechanism from the growth factor inhibitors. The cytotoxic effect of the taxanes is caused by interference with tubulin, which prevents normal mitosis (cell division). This is the main effect seen at the high doses of the taxanes used in cancer chemotherapy. A secondary effect is disruption of newly formed blood vessels, since the cells of the new vascular endothelium depend on tubulin to maintain their shape. However, this effect is normally seen only at doses too low to be cytotoxic. Any synergy between DMXAA and the taxanes is thought to be a result of the targeting of different parts of the tumour, as described above, rather than due to its anti-angiogenic properties.

Other agents have also been shown to enhance the activity of DMXAA in xenograft studies. Although the exact mechanism of action of DMXAA is not understood, it is believed to cause upregulation of various cytokines, and compounds with similar activity appear to enhance its effectiveness. These include tumour necrosis factor stimulating compounds and immunomodulatory compounds such as intracellular adhesion molecules (ICAMs).

Diclofenac, an NSAID that has been shown to enhance the anti-tumour activity of DMXAA, is believed to affect the PK of DMXAA via competition for metabolic pathways. At a concentration of 100µM, diclofenac has been shown to significantly inhibit glucuronidation (>70%) and 6-methylhydroxylation (>54%) of DMXAA in mouse and human liver microsomes. *In vivo*, diclofenac (100mg/kg i.p.) has been shown to result in a 24% and 31% increase in the plasma DMXAA AUC (area under the plasma concentration-time curve) and a threefold increase in $T_{1/2}$ ($P<0.05$) in male and female mice respectively (Zhou et al (2001) Cancer Chemother Pharmacol 47 319-326). Other NSAIDs have been shown to have a similar effect.

Similarly to diclofenac, thalidomide, which is approved for erythema nodosum leprosum (ENL), seems to enhance the activity of DMXAA. It competes for glucuronidation, prolonging DMXAA's presence at therapeutic levels in tumour tissue. Thalidomide increases the AUC of DMXAA by 1.8 times in plasma, liver and spleen and by three times in tumour (Kestell et al. Cancer Chemother Pharmacol. 2000;46(2):135-41). Thalidomide is known to have anti-angiogenic effects, but these are not believed to be responsible for its synergy with DMXAA. It would not be expected that combining with vascular endothelial growth factor binder would have a similar effect to that of thalidomide on the effectiveness of DMXAA.

Previous vascular endothelial growth factor binder combination studies

Clinical evidence teaches away from combining different types of vascular targeting agents. It has been shown that Avastin™ does not have a synergistic effect when used in combination with thalidomide, an angiogenesis inhibitor, in metastatic renal cell carcinoma (Elaraj et al. J Immunother. 2004 Jul-Aug ; 27(4) : 259-64).


Progression-free survival was the same in patients treated with Avastin™ alone or Avastin™ combined with thalidomide. In its approved indication, colorectal cancer, Avastin™ is used in combination with 5-FU (5-fluorouracil), which does not have anti-angiogenic properties. Avastin™ has also been shown to improve median survival in breast and lung cancer patients when combined with paclitaxel. Although paclitaxel does have some anti-angiogenic properties, its primary mechanism of action in the high doses in which it is used for cancer treatment is as a cytotoxic, as described above. Therefore, this would not suggest that DMXAA would have a

similar synergy with Avastin®, since DMXAA is very unlike paclitaxel in its mechanism of action and is not a cytotoxic.

Description of the invention

In a first aspect, the present invention provides a method for modulating neoplastic growth, which comprises administering to a mammal, including a human, in need of treatment a compound of formula I:

Formula I:




wherein:

- (a) R_4 and R_5 together with the carbon atoms to which they are joined, form a 6-membered aromatic ring having a substituent $-R_3$ and a radical $-(B)-COOH$ where B is a linear or branched substituted or unsubstituted C_1-C_6 alkylene radical, which is saturated or ethylenically unsaturated, and wherein R_1 , R_2 and R_3 are each independently selected from the group consisting of H, C_1-C_6 alkyl, halogen, CF_3 , CN, NO_2 , NH_2 , OH, OR, $NHCOR$, $NHSO_2R$, SR, SO_2R or NHR, wherein each R is independently C_1-C_6 alkyl optionally substituted with one or more substituents selected from hydroxy, amino and methoxy; or
- (b) one of R_4 and R_5 is H or a phenyl radical, and the other of R_4 and R_5 is H or a phenyl radical which may optionally be substituted, thenyl, furyl, naphthyl, a C_1-C_6 alkyl, cycloalkyl, or aralkyl radical; R_1 is H or a C_1-C_6 alkyl or C_1-C_6 alkoxy radical; R_2 is the radical $-(B)-COOH$ where B is a linear or branched substituted or unsubstituted C_1-C_6 alkylene radical, which is saturated or ethylenically unsaturated,

or a pharmaceutically acceptable salt, ester or prodrug thereof and concomitantly or sequentially administering a vascular endothelial growth factor binder.

Where (B) in the radical $-(B)-COOH$ is a substituted C_1-C_6 alkyl radical, the substituents may be alkyl, for example methyl, ethyl, propyl or isopropyl, or halide such as fluoro, chloro or bromo groups. A particularly preferred substituent is methyl.


In one embodiment of the first aspect of the invention, the compound of the formula (I) as defined above is a compound of the formula (II),



where R_1 , R_4 , R_5 and B are as defined above for formula (I) in part (b).

In a further embodiment of the first aspect of the invention, the compound of formula (I) as defined above is a compound of the formula (III)

Formula (III)



wherein R_1 , R_2 and R_3 are each independently selected from the group consisting of H, C_1 - C_6 alkyl, halogen, CF_3 , CN, NO_2 , NH_2 , OH, OR, $NHCOR$, $NHSO_2R$, SR, SO_2R or NHR , wherein each R is independently C_1 - C_6 alkyl optionally substituted with one or more substituents selected from hydroxy, amino and methoxy;


wherein B is as defined for formula (I) above;

and wherein in each of the carbocyclic aromatic rings in formula (I), up to two of the methine ($-CH=$) groups may be replaced by an aza ($-N=$) group;

and wherein any two of R_1 , R_2 and R_3 may additionally together represent the group $-CH=CH-CH=CH-$, such that this group, together with the carbon or nitrogen atoms to which it is attached, forms a fused 6 membered aromatic ring.

For example, the compound of formula (III) may be a compound of the formula (IV):

Formula (IV)




wherein R, R_1 , R_2 and R_3 are as defined for formula (III).

In one embodiment of the compound of formula (IV), R_2 is H, one of R_1 and R_3 is selected from the group consisting of C_1 - C_6 alkyl, halogen, CF_3 , CN, NO_2 , NH_2 , OH, OR, $NHCOR$, $NHSO_2R$, SR, SO_2R or NHR , wherein each R is independently C_1 - C_6 alkyl optionally substituted with one or more substituents selected from hydroxy, amino and methoxy, and the other of R_1 and R_3 is H.

In one embodiment, in the compound of formula (I) R_4 is H or a phenyl radical, R_5 is H or a phenyl radical which may optionally be substituted, thenyl, furyl, naphthyl, a C_1 - C_6 alkyl, cycloalkyl, or aralkyl radical; R_1 is H or a C_1 - C_6 alkyl or C_1 - C_6 alkoxy radical; R_2 is radical $-(B)-COOH$ where B is a linear or branched substituted or unsubstituted C_1 - C_6 alkylene radical, which is saturated or ethylenically unsaturated.

For example, the compound of formula (IV) may be a compound of the formula (V):

Formula (V)



wherein R, R_1 , R_2 and R_3 are as defined for formula IV.

The compound of formula (V) may be, for example, 5,6-dimethylxanthhenone-4-acetic acid (DMXAA).

In another aspect, the present invention provides the use of a vascular endothelial growth factor binder for the manufacture of a medicament (e.g. a unit dose of the medicament), for simultaneous, separate or sequential administration with a compound of formula (I) as defined above or a pharmaceutically acceptable salt, ester or prodrug thereof (e.g. a unit dose of the compound of formula (I) as defined above or a pharmaceutically acceptable salt, ester or prodrug thereof), for the modulation of neoplastic growth.

In a further aspect, the invention provides the use of a compound of formula (I) as defined above or a pharmaceutically acceptable salt or ester thereof for the manufacture of a medicament (e.g. a unit dose of the medicament), for simultaneous, separate or sequential administration with a vascular endothelial growth factor binder

(e.g. a unit dose of the vascular endothelial growth factor binder), for the modulation of neoplastic growth.

According to one aspect, the neoplastic growth is a tumour and/or a cancer.

In a further aspect, the cancer is one or more of ovarian, prostate, lung, colorectal, breast, pancreatic and renal cancer.

In a further aspect, there is provided a pharmaceutical formulation (e.g. in a unit dose) comprising a combination of a compound of formula (I) as defined above or a pharmaceutically acceptable salt or ester or prodrug thereof (e.g. in a unit dose) and a vascular endothelial growth factor binder (e.g. in a unit dose).

In one embodiment there is provided a compound according to formula I or a pharmaceutically acceptable salt, ester or prodrug thereof and a vascular endothelial growth factor binder for use (in combination) as a medicament for the modification of neoplastic growth.

Furthermore, the invention also provides a kit comprising in combination for simultaneous, separate or sequential use in modulating neoplastic growth, a compound of formula (I) as defined above or a pharmaceutically acceptable salt or ester or prodrug thereof and a vascular endothelial growth factor binder.

The compound of formula (I) as defined above or pharmaceutically acceptable salt or ester or prodrug thereof and the vascular endothelial growth factor binder may be administered sequentially or concomitantly. For example, the compound of formula (I) as defined above or pharmaceutically acceptable salt, ester or prodrug thereof and the vascular endothelial growth factor binder may be administered concomitantly.

In one embodiment, the pharmaceutically acceptable salt is a sodium salt.

The compound of formula (I) as defined above or pharmaceutically acceptable salt, ester or prodrug thereof and the vascular endothelial growth factor binder may be administered simultaneously, separately or sequentially.

In one embodiment, the vascular endothelial growth factor binder is a monoclonal antibody.

In a further embodiment, vascular endothelial growth factor binder (VEGF) is AvastinTM (bevacizumab).

The amount of a combination of a compound of formula (I) as defined above or pharmaceutically acceptable salt, ester or prodrug thereof and a vascular endothelial growth factor binder required to be effective as a modulator of neoplastic growth will, of course vary and is ultimately at the discretion of the medical practitioner. The factors to be considered include the route of administration and nature of the formulation, the mammal's bodyweight, age and general condition and the nature and severity of the disease to be treated.

A suitable effective dose of a compound of formula (I) as defined above, or a pharmaceutically acceptable salt, ester or prodrug thereof, for administration, concomitantly or sequentially, with a vascular endothelial growth factor binder, for the treatment of cancer is in the range of 600 to 4900mg/m². For example from 2500 to 4000 mg/m², for example from 1200 to 3500mg/m², for example from 2000 to 3000 mg/m², for example from 1200 to 2500 mg/m², for example from 2500 to 3500 mg/m², for example from 2250 to 2750 mg/m².

A suitable effective dose of vascular endothelial growth factor binder, for administration concomitantly or sequentially with a compound of formula (I) as defined above or pharmaceutically acceptable salt, ester or prodrug thereof for the treatment of cancer is in the range of 1-10mg/kg, for example about 5mg/kg.

A compound of formula (I) as defined above or pharmaceutically acceptable salt, ester or prodrug thereof and the vascular endothelial growth factor binder may be administered in any suitable form, for example in the form of a pharmaceutical formulation.

Pharmaceutical formulations comprise the active ingredients (that is, the combination of a compound of formula (I) as defined above or pharmaceutically acceptable salt, ester or prodrug thereof and the vascular endothelial growth factor binder, for example together with one or more pharmaceutically acceptable carriers therefor and optionally other therapeutic and/or prophylactic ingredients. The carrier(s) must be acceptable in the sense of being compatible with the other ingredients in the formulation and not deleterious to the recipient thereof.

Accordingly, the present invention provides a pharmaceutical formulation comprising a combination of a compound of formula (I) as defined above or pharmaceutically acceptable salt, ester or prodrug thereof (e.g. a unit dose of a compound of formula (I) as defined above or pharmaceutically acceptable salt, ester or prodrug thereof) and a vascular endothelial growth factor binder (e.g. a unit dose of the vascular endothelial growth factor binder), for example in association with one or more pharmaceutically acceptable carriers therefor.

The invention further provides a process for the preparation of a pharmaceutical formulation which process comprises bringing into association a combination of a compound of formula (I) as defined above or a pharmaceutically acceptable salt, ester or prodrug thereof (e.g. a unit dose of a compound of formula (I) as defined above or pharmaceutically acceptable salt, ester or prodrug thereof) and a vascular endothelial growth factor binder (e.g. a unit dose of the vascular endothelial growth factor binder) optionally together with one or more pharmaceutically acceptable carriers therefor in. For example, the pharmaceutical formulation may be in a unit dose.

The pharmaceutical formulation may be delivered intravenously. The pharmaceutical formulation for intravenous administration may be used in the form of sterile aqueous solutions or in an oleaginous vehicle which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood. The aqueous solutions may be buffered (e.g. to a pH from 3 to 9), if necessary.

As used herein, the term "prodrug" includes entities that have certain protected group(s) and which may not possess pharmacological activity as such, but may, in

certain instances, be administered (such as orally or parenterally) and thereafter metabolised in the body to form the agent which are pharmacologically active.

Examples

Xenografts for human lung and colorectal cancers are set-up in groups of nude, athymic mice. The cell lines selected are HT29 (ATCC Number HTB-38), a colorectal adenocarcinoma, and A549 (ATCC Number CCL-185), a lung carcinoma.

The A549 and HT29 cell lines are selected as DMXAA has previously been shown to be effective in these cell lines when used in combination with paclitaxel or 5-FU in xenograft studies. In addition, Avastin™ is currently approved for treatment of colorectal cancer in combination with 5-FU and approval is being sought for use on breast and non-small cell lung carcinoma.

Group	Cell line	Treatment	Dose level (mg/kg)	No. of mice
1	A549	Untreated control	-	10
2	A549	DMXAA	21	10
3	A549	Avastin™	5	10
4	A549	DMXAA + Avastin™	21 & 5	10
5	HT29	Untreated control	-	10
6	HT29	DMXAA	21	10
7	HT29	Avastin™	5	10
8	HT29	DMXAA + Avastin™	21 & 5	10


DMXAA has been given previously using a day (D) 0, 4 and 8 schedule when used in combination with paclitaxel or docetaxel. For this study, DMXAA is given twice in each of Weeks 1 and 4 of the study. Avastin™ is given twice weekly for four weeks.

Xenografts are measured two or three times per week and their absolute volume recorded; xenograft tumour volume relative to that recorded on Day 0 (V_0) is then calculated. The time taken to reach a relative tumour volume of $3 \times V_0$ is used as a surrogate marker for survival.

CLAIMS

1. A method for modulating neoplastic growth, which comprises administering to a mammal, including a human, in need of treatment a compound of formula (I):

Formula (I)




wherein:

- (a) R₄ and R₅ together with the carbon atoms to which they are joined, form a 6-membered aromatic ring having a substituent -R₃ and a radical -(B)-COOH where B is a linear or branched substituted or unsubstituted C₁-C₆ alkylene radical, which is saturated or ethylenically unsaturated, and wherein R₁, R₂ and R₃ are each independently selected from the group consisting of H, C₁-C₆ alkyl, halogen, CF₃, CN, NO₂, NH₂, OH, OR, NHCOR, NHSO₂R, SR, SO₂R or NHR, wherein each R is independently C₁-C₆ alkyl optionally substituted with one or more substituents selected from hydroxy, amino and methoxy; or
- (b) one of R₄ and R₅ is H or a phenyl radical, and the other of R₄ and R₅ is H or a phenyl radical which may optionally be substituted, thenyl, furyl, naphthyl, a C₁-C₆ alkyl, cycloalkyl, or aralkyl radical; R₁ is H or a C₁-C₆ alkyl or C₁-C₆ alkoxy radical; R₂ is the radical -(B)-COOH where B is a linear or branched substituted or unsubstituted C₁-C₆ alkylene radical, which is saturated or ethylenically unsaturated,

or a pharmaceutically acceptable salt, ester or prodrug thereof and concomitantly or sequentially administering a vascular endothelial growth factor binder,


2. The method according to claim 1 wherein the compound of Formula (I) is a compound of Formula (II):

Formula (II)



wherein R_1 , R_4 , R_5 and B are as defined for formula (I) in claim 1 part (b).

3. The method according to claim 1 wherein the compound of Formula (I) is a compound of Formula (III):



Formula (III)

wherein R_1 , R_2 and R_3 are each independently selected from the group consisting of H, C_1 - C_6 alkyl, halogen, CF_3 , CN, NO_2 , NH_2 , OH, OR, $NHCOR$, $NHSO_2R$, SR, SO_2R or NHR , wherein each R is independently C_1 - C_6 alkyl

optionally substituted with one or more substituents selected from hydroxy, amino and methoxy;


wherein B is as defined for formula (I) in claim 1;

and wherein in each of the carbocyclic aromatic rings in formula (I), up to two of the methine (-CH=) groups may be replaced by an aza (-N=) group;

and wherein any two of R₁, R₂ and R₃ may additionally together represent the group -CH=CH-CH=CH-, such that this group, together with the carbon or nitrogen atoms to which it is attached, forms a fused 6 membered aromatic ring.

4. The method according to claim 3, wherein the compound of Formula (III) is a compound of Formula (IV):


Formula (IV)



wherein R, R₁, R₂ and R₃ are as defined for formula (III) in claim 3.

5. The method according to claim 4 wherein the compound of Formula (IV) is a compound of formula (V):

17



Formula (V)

wherein R, R₁, R₂ and R₃ are as defined for formula IV in claim 4.

6. The method according to claim 1, wherein the compound of Formula (I) is DMXAA or a pharmaceutically acceptable salt, ester or prodrug thereof.
7. A method according to any of claims 1 to 6 wherein the compound of formula (I) or a pharmaceutically acceptable salt, ester or prodrug thereof and the vascular endothelial growth factor binder are administered concomitantly.
8. A method according to any one of the claims 1 to 6 wherein the compound of formula (I) or pharmaceutically acceptable salt, ester or prodrug thereof and the vascular endothelial growth factor binder are administered sequentially.
9. The method according to any one of the preceding claims wherein the vascular endothelial growth factor binder is a monoclonal antibody.
10. The method according to claim 9 wherein the vascular endothelial growth factor binder is AvastinTM (bevacizumab).
11. The method according to any one of the preceding claims wherein the method further comprises modulation of neoplastic growth in one of more of ovarian, prostate, lung, colorectal, pancreatic, breast and renal cancer.

12. Use of the compound according to any one of claims 1 to 6 or a pharmaceutically acceptable salt, ester or prodrug thereof for simultaneous, separate or sequential administration with a vascular endothelial growth factor binder, for the modulation of neoplastic growth.
13. Use according to claim 12 wherein the vascular endothelial growth factor binder is a monoclonal antibody.
14. Use according to claim 13 wherein the vascular endothelial growth factor is Avastin™ (bevacizumab).
15. Use according to claim 12, 13 or 14 wherein the compound is DMXAA or a pharmaceutically acceptable salt, ester or prodrug thereof.
16. Use according to any one of claims 12 to 15 wherein the modulation of neoplastic growth is in one of more of ovarian, prostate, lung, colorectal, pancreatic, breast and renal cancer.
17. Use of a vascular endothelial growth factor binder for the manufacture of a medicament, for simultaneous, separate or sequential administration with a compound according to any one of claims 1 to 6 or a pharmaceutically acceptable salt, ester or prodrug thereof, for the modulation of neoplastic growth.
18. Use of the compound of formula (I) according to any one of claims 1 to 6 or a pharmaceutically acceptable salt, ester or prodrug thereof for the manufacture of a medicament, for simultaneous, separate or sequential administration with a vascular endothelial growth factor binder, for the modulation of neoplastic growth.
19. Use according to claim 17 or claim 18 wherein the vascular endothelial growth factor binder is bevacizumab.
20. Use according to any one of claims 17 to 19 wherein compound of formula (I) is DMXAA or a pharmaceutically acceptable salt, ester or prodrug thereof.

21. A pharmaceutical formulation comprising a combination of a compound according to any one of claims 1 to 6 or a pharmaceutically acceptable salt, ester or prodrug thereof and a vascular endothelial growth factor binder.
22. The pharmaceutical formulation of claim 21 wherein the pharmaceutical formulation further comprises a pharmaceutically acceptable carrier.
23. A pharmaceutical formulation according to claim 21 or claim 22 wherein the formulation is adapted for intravenous administration.
24. A pharmaceutical formulation according to any one of claims 21 to 23 wherein the vascular endothelial growth factor binder bevacizumab.
25. A pharmaceutical formulation according to any one of claims 21 to 24 wherein the compound of formula (I) is DMXAA or a pharmaceutically acceptable salt, ester or prodrug thereof.
26. A compound according to any one of claims 1 to 6 and vascular endothelial growth factor binder for modulating neoplastic growth.
27. The compounds according to claim 26 wherein the vascular endothelial growth factor binder is bevacizumab.
28. The compounds according to claim 26 or claim 27 wherein the compound of formula (I) is DMXAA or a pharmaceutically acceptable salt, ester or prodrug thereof.
29. A process for the preparation of a pharmaceutical formulation which process comprises bringing into association a compound according to any one of claims 1 to 6 or a pharmaceutically acceptable salt, ester or prodrug thereof and a vascular endothelial growth factor binder.
30. The process according to claim 29 wherein the compound according to any one of claims 1 to 6 or a pharmaceutically acceptable salt, ester or prodrug thereof and

the vascular endothelial growth factor binder are brought into association with one or more pharmaceutically acceptable carriers therefor.

31. The process according to claim 29 or 30 wherein the vascular endothelial growth factor binder is bevacizumab.

32. The process according to any of claims 29 to 31 wherein the compound of formula (I) is DMXAA or a pharmaceutically acceptable salt, ester or prodrug thereof.

33. A kit comprising in combination for simultaneous, separate or sequential use in modulating neoplastic growth, a compound according to any one of claims 1 to 6 or a pharmaceutically acceptable salt, ester or prodrug thereof and a vascular endothelial growth factor binder.

34. The kit according to claim 33 wherein the growth factor inhibitor is bevacizumab.

35. The kit according to claim 32 or 33 wherein the compound of formula (I) is DMXAA or a pharmaceutically acceptable salt, ester or prodrug thereof.

ABSTRACT**COMBINATIONS FOR THE TREATMENT OF CANCER**

The present invention relates to combinations of compounds such as compounds of the xanthenone acetic acid class such as 5,6-dimethylxanthenone-4-acetic acid (DMXAA) and vascular endothelial growth factor binders, in particular the monoclonal antibody Avastin™ (bevacizumab). More particularly, the invention is concerned with the use of such combinations in the treatment of cancer and pharmaceutical formulations containing such combinations.

